

Marquis (15% w/w ponazuril) Antiprotozoal Oral Paste Indications

Marquis (ponazuril) is indicated for the treatment of equine protozoal myeloencephalitis (EPM) caused by *Sarcocystis neurona*.

DOSAGE: Marquis (15% w/w ponazuril) is to be used at a dose of 5 mg/kg (2.27 mg/lb) body weight once daily for a period of 28 days

ADMINISTRATION:

Paste syringe assembly:


Before administration, the syringe barrel and plunger require

assembly. Ensure plunger is clean and dry.

Step 1. End cap **must** be on syringe barrel when inserting plunger.

Step 2. Carefully insert reusable plunger into base of syringe barrel until it snaps into place, then remove end cap and gently apply pressure to the plunger until paste is seen at the tip of the syringe barrel.

Step 3. Return end cap to tip of paste syringe.



Administering Marquis (ponazuril) to the horse:

NOTE: The paste syringe is a multi-dose package.

Ensure that

the correct dose is administered with each use.

Step 1. Remove end cap and gently apply pressure to the plunger

until paste is seen at the tip of the syringe barrel. Return end cap to

tip of paste syringe.

Step 2. Determine weight of horse and insure the horse's mouth

contains no feed.

Step 3. To measure dose, dosage ring collar and barrel

should be flush. Hold plunger and rotate dosage ring with the other hand to the weight of the horse.

Step 4. Remove end cap from tip of syringe barrel.

Step 5. The selected dose of paste should be deposited onto the

back and top of the horse's tongue. Introduce tip of paste syringe into

the side of the horse's mouth at the space between the front (incisor)

and back (molar) teeth. Deposit paste on the horse's tongue by

depressing the plunger of the syringe as far as the dose ring permits. Remove tip of syringe from horse's mouth.

Step 6. To aid swallowing of paste, immediately raise horse's head for

a few seconds after dosing.

Step 7. Clean the tip of the syringe with a clean disposable towel and

return end cap to tip of syringe barrel.

Step 8. For the next daily dose, repeat steps 1-7.

