

### Endo-parasitic drugs

Drug	Use	Animal	Route of administration	Dosage	Warning	Withdrawal period
Equitak (broad spectrum wormer for horses)	Treatment and control of roundworms, tapeworms and bots in horses.  Contains the unique 3-in-1 combination of active ingredients, Abamectin, Oxfendazole and Praziquantel.	Horses	Oral administration	1g of Equitak Excel per 20kg bodyweight.  Repeat doses should be given every 6 - 8 weeks in young and more susceptible horses.  Mature horses only require treatment 3 - 4 times per year.	Dewormer should be properly rotated to prevent resistance from parasites.	Withdraw for 28 to 30 days before slaughter for human consumption
Chanaverm	A broad spectrum anthelmintic for the treatment and control of gastro-intestinal and pulmonary nematode infections in cattle and sheep.	Cattle sheep	Oral administration	Cattle: 1 mL product/10 kg body weight.  Sheep: 0.5 mL product/5 kg body weight.	Underdosing and repeated usage of anthelmintics over an extended period of time can cause resistance build up	Meat and offal: 18 days.  Not for use in animals producing milk for human consumption
Amprolium 250	Antiprotozoal drug use to	Cattle Swine	Oral Administration	For coccidiosis in cattle:	A withdrawal period has not been	Withdrawal time for cattle (meat): 24 hours before slaughter

<p>Ancoban [Amprolium 20%]</p>	<p>antagonizes thiamine in parasites and use for treatment of coccidiosis</p>	<p>Goat Sheep</p>		<p>Treatment- 10 mg/kg PO for 5 days; 5 mg/kg for 21 days for Prophylaxis.</p> <p>For coccidiosis in swine: Treatment- 25 – 65 mg/kg PO once or twice daily for 3 – 4 Days or 100 mg/kg/day in food or water.</p> <p>For coccidiosis In sheep and goats: 55 mg/kg daily PO for 19 days</p>	<p>established for this product in preruminating calves. So do not use in calves to be processed for veal.</p> <p>Do not administer to debilitated animals.</p>	
<p>Imizol Injection</p>	<p>Antiprotozoal drug used in animals to treat intracellular tickborne pathogens. It has been used to treat <i>Babesia</i> infections.</p>	<p>Horses Cattle Sheep</p>	<p>Intramuscular Subcutaneous</p>	<p>Horses: For treatment of equine piroplasmosis- 2.2 mg/kg IM will generally allow clinical signs to subside. To eliminate <i>B.</i></p>	<p>Do not use imidocarb in patients exposed to cholinesterase-inhibiting drugs, pesticides, or chemicals.</p>	<p>28 days for meat and 3 days for milk.</p>

				<p><i>caballi</i> inject 2 mg/kg IM once a day for 2 days.</p> <p>Sheep: For treatment of babesiosis- 1.2 mg/kg IM; repeat in 10 – 14 days.</p> <p>Treatment of Anaplasmosis &amp; mixed infections due to Anaplasma &amp; Babesia in cattle: 2.5ml per 100kg bodyweight.</p>		
--	--	--	--	---	--	--