

# APPLICATION TECHNIQUES OF LOCAL BLOCK ANAESTHESIA IN RUMINANTS

LOCAL anaesthesia is a commonly used technique in veterinary practice and many surgical procedures in ruminants can be carried out under local blocks.

It is used routinely for procedures such as dehorning, as well as for surgeries such as caesarean section and correction of abomasal displacements. Sedation may be employed as an adjunct (see the first article of this series, VT37.03) but its use depends on the animal's species, temperament and health. Pre-emptive local anaesthesia can be used in combination with general anaesthesia to reduce the required dose of general anaesthetic product and minimise the potential cardiopulmonary depression that may accompany it and also assist in a quicker recovery.

Current licensing limits the drugs that may be used for induction of local anaesthesia in ruminants. The only local anaesthetic currently licensed for use in food-producing species in the UK is procaine. Lignocaine (lidocaine in North America) had become extremely widely used, and its dose rates are widely reported in the literature; however, its use

## GAYLE D HALLOWELL

MA, VetMB, CertVA, DipACVIM, MRCVS, and

## TIMOTHY J POTTER

BVetMed, MRCVS

continue their discussions on anaesthesia in ruminants by looking at use of local blocks

in the UK is now off label. Compared with procaine, lignocaine has a far shorter period of onset, as well as having a more intense effect and a longer duration of action. Procaine's poorer spread through tissues necessitates more accurate deposition within the tissues than lignocaine.

Toxicity associated with local anaesthetics is a concern – especially in small ruminants. Excessive plasma concentrations of local anaesthetics can block cardiac sodium channels and depress conduction of impulses through the heart. Further cardiac toxicity is caused by the inhibition of cyclic adenosine monophosphate (cAMP) production and the local anaesthetic action on calcium and potassium channels.

Clinical signs of overdose are

initial sedation, followed with increasing dosage by twitching convulsions, coma and death. The toxic dose of lignocaine is 8mg/kg and for procaine 20mg/kg. Such concentrations can be inadvertently reached, espe-

cially in smaller animals when using two per cent solutions for infiltration blocks.

### Blocks for standing laparotomy

No fewer than six techniques for inducing anaesthesia of the paralumbar fossa and abdominal wall in standing ruminants have been described:

- infiltration (for example, line block or inverted L);
- proximal paravertebral thoracolumbar;
- distal paravertebral thoracolumbar;

- segmental dorsolumbar epidural;
- continuous lumbar segmental epidural; and
- thoracolumbar subarachnoid anaesthesia.

The choice of method is down to operator preference and Table 1 summarises the advantages and disadvantages of the different techniques. It is beyond the scope of this article to provide a detailed description of all the available techniques. A description of the proximal vertebral is included as an example, as this is the authors' method of choice.

### ● Proximal paravertebral block

This technique involves anaesthetising the spinal nerves as they emerge from their intervertebral foramina. It is often sufficient to anaesthetise just the dorsal and ventral branches of the last thoracic (T13) and the first two lumbar (L1 and L2) nerves.

Some procedures, such as caesarean section, require inclusion of the third and fourth lumbar (L3 and L4) nerves. It should be remembered that inclusion of L3 and L4 may result in ataxia as these nerve supply motor fibres

**TABLE 1.** Summary of local anaesthetic techniques employed for laparotomy in ruminants

Technique	Advantages	Disadvantages
Line block	Easy.	Oedema and haematoma along the incision site may interfere with healing. Incomplete analgesia and muscle relaxation of the deeper layers of the abdominal wall. Potential toxicity after injection of significant amounts of analgesia. Large volume.
Inverted L block	Easy. Deposition of anaesthetic away from the incision.	Incomplete analgesia and muscle relaxation of the deeper layers of the abdominal wall. Potential toxicity after injection of significant amounts of analgesia. Large volume.
Proximal paravertebral block	Small volume. Wide and uniform area of analgesia and muscle relaxation. Increased intestinal tone and motility. Absence of local anaesthetic from the wound edges.	Technical difficulty. Arching up of the spine due to paralysis of the back muscles. Risk of penetrating vital structures.
Distal paravertebral block	No risk of penetrating a major blood vessel. Lack of scoliosis. Minimal weakness in pelvic limbs and ataxia.	Technical difficulty. Larger dose than proximal paravertebral. Variable efficiency.
Thoracolumbar subarachnoid anaesthesia	Rapid onset. Small volume. Catheter used provides a port for repeated boluses of anaesthetic.	Requirement for specific equipment. Risk of trauma to conus medullaris. Ataxia or recumbency after overdose. Risk of meningitis.
Segmental dorsolumbar epidural	One injection site. Small volume. Uniform anaesthesia and relaxation of skin musculature and peritoneum.	Technically difficult. Risk of trauma to spinal cord and venous sinuses. Potential loss of motor control of pelvic limbs. Potential physiological disturbances (such as bradycardia and hypotension).
Continuous lumbar segmental epidural	Small doses can be administered so the extent of anaesthesia can be readily controlled.	Technically difficult. Requires specialised equipment. Increased frequency of post-anaesthetic myositis

PRAC-TIC WITH PYRIPROLE.  
FREEDOM FROM  
FLEAS AND TICKS



EFFECTIVE EVEN  
AFTER BATHING OR  
WATER IMMERSION\*

to the ischial femoral nerves.

The first step is to identify the transverse processes by careful palpation. Start caudally by identifying the process of L5, which lies just in front of the tuber coxae, then work cranially, ending up on L1, which is much smaller and often harder to palpate. Skin overlying the dorsal aspects of the transverse processes is clipped and scrubbed. A small quantity of local anaesthetic is injected just under the skin on the dorsal aspect of the transverse processes of L1, L2 and L3, 5cm from midline.

To anaesthetise L1 a 19-gauge, 3.5in spinal needle is inserted through the desensitised skin on the cranial edge of L2 and advanced ventrally (Figure 1). If the needle impacts the transverse process it is "walked off" the cranial aspect and passed ventrally through the transverse

ligament and fascia. The penetration of the intertransverse fascia can normally be felt – 15ml of local anaesthetic is then introduced. The needle is then withdrawn 1-2cm so that the needle tip lies above the fascia and 5ml of anaesthetic is injected against slight resistance to desensitise the dorsal branch of L1. To anaesthetise T13 and L2 the needle is inserted on cranial aspect of the transverse processes of L1 and L3 respectively and the method is repeated as for L1.

**Caudal epidural anaesthesia**

This is an inexpensive technique that does not require any specialised equipment. It is routinely used in cattle, sheep and goats – most frequently for obstetrical procedures. The aim is to desensitise the last three pairs of sacral nerves within the spinal

column without compromising the motor function of the pelvic limbs. Needle placement is either at the sacrococcygeal (S5-Co1) interspace or, more commonly, at the first coccygeal (Co1-Co2) interspace (Figure 2). The first coccygeal joint is easily identified by moving the tail up and down and palpating the movement between the Co1 and Co2 vertebrae. In contrast, it can be harder to identify the sacrococcygeal joint, as it is an immovable joint and may become calcified in cattle more than eight years of age.

The skin overlying the point of needle insertion should be clipped, scrubbed and disinfected. An 18-gauge, 1.5in needle is normally sufficient, although in large animals or those carrying excessive condition, a longer needle may be required.

The point of the needle

is placed in the centre of the depression and advanced cranially and ventrally at an angle of 15° to the vertical. The needle is advanced until it contacts the ventral floor of the vertebral canal. The needle is then withdrawn approximately 0.5cm into the epidural space. Correct placement can be identified by placing a couple of drops of anaesthetic in the hub of the needle; sub-atmospheric pressure in the epidural space should lead to them being aspirated (hanging drop technique). Local anaesthetic can then be administered. No resistance should be encountered to the injection.

In cattle, anaesthesia extending cranially from the middle of the sacrum and ventrally over the perineum to the inner aspect of the thigh can be achieved with a dose of 1ml of two per cent lignocaine per 100kg bodyweight<sup>1</sup>. At

Figure 1. Careful palpation of the transverse processes allows accurate needle placement for a proximal paravertebral block.



an appropriate dose range, pelvic viscera and genitalia are anaesthetised and abdominal contractions are abolished while leaving uterine motility and the locomotor function of the hindlimbs unaffected. Maximal effect is seen 10 to 15 minutes after administration and lasts for around one hour. Lack of anaesthetic effect within 10 minutes usually suggests that the injection was made outside of the epidural space.

Complications are rare but can include postural instability and/or recumbency and haemorrhage due to puncture of a venous sinus. In cases where asepsis is not observed, infection may lead to a discharging tract or a permanently paralysed tail.

Caudal epidural anaesthesia in sheep and goats can be performed in a similar manner to cattle – again ensuring asepsis

is observed. The maximum recommended dose rate is 1ml of two per cent lignocaine per 50kg bodyweight.

The use of regional anaesthetic techniques in farm animal practice is often chosen above general anaesthesia due to its ease of administration, low cost and reduced incidence of complications. In addition to providing a means of anaesthesia, regional anaesthetic techniques should not be overlooked as a component of multimodal analgesia.

The use of local anaesthetics, either alone or in combination with other analgesics, can provide a useful tool for providing effective analgesia in ruminants.

**Reference**

1. Skarda R T (1996). Local and regional anesthesia in ruminants and swine, *Vet Clin North Am Food Anim Pract* 12: 579-626. ■




Figure 2. The red line indicates the needle placement between the Co1 and Co2 vertebrae.

When you prescribe Prac-tic, you're prescribing a treatment that kills 100% of fleas<sup>†</sup> in 24 hours so completely eliminates egg production and breaks the flea lifecycle, kills 99% of ticks<sup>†</sup> in 48 hours, and offers dogs a full month's protection that is effective even after bathing or water immersion.\*

**Prac-tic**<sup>®</sup>  
find your freedom

For further information contact your vet, or in the UK the Practice support line on 0800 845100, or write to Novartis Animal Health UK Ltd, Frimley Business Park, Frimley, Camberley, Surrey GU16 7SR. For further information in Ireland telephone 051 377201, or write to Novartis Animal Health Ireland Ltd, Industrial Park, Cork Road, Waterford, Eire. Prac-tic (POM) in the UK, and (POM) in Ireland contains pyriprole. Prac-tic<sup>®</sup> is a registered trademark of Novartis AG, Basel, Switzerland. © 2008 Novartis Animal Health Inc, Basel, Switzerland. \*NAH 04-0015, NAH 03-0005, CH001/04-009. Dogs should not be bathed 48 hours, before, or 24 hours after treatment. <sup>†</sup>NAH CH001/03-003 (Tick spp: R.sanguineus) <sup>††</sup>Subject to MA approval in the country of use.



**GAYLE HALLOWELL** graduated from the University of Cambridge and completed an equine internship and a residency in equine medicine and critical care at the Royal Veterinary College, London. She is a temporary lecturer in equine medicine at the RVC and has started a PhD training programme, funded by the Horserace Betting Levy Board, at the University of Nottingham School of Veterinary Medicine and Science.

**TIM POTTER** graduated from the Royal Veterinary College and completed a residency in dairy herd medicine and surgery based jointly at the RVC and the Shepton Veterinary Group in Somerset. He is currently undertaking a PhD training programme, funded by DEFRA, examining the design of dosing regimes to minimise the development of antimicrobial resistance in calf pneumonia.

